LBRC Report Form

- 1. English and Scientific names: Hooded Oriole (Icterus cucullatus)
- 2. Number of individuals, sexes, ages, general plumage (e.g., 2 in alternate plumage):
- 1, f-type plumage.
- 3. Parish:

Jefferson Parish

4. Specific Locality:

Grand Isle, LA 179 Landry Ln. in Turks Caps next to bed and breakfast

- 5. Date(s) when observed: Feb 25th, 26th, 27th, 28th, 29th, Mar 1st
- 6. Time(s) of day when observed: Early morning and afternoon
- 7. Reporting observer: Jennie Leonard
- 8. Other observers accompanying reporter who also identified the bird(s): Shane Schopinsky (partner)
- 9. Other observers who independently identified the bird(s): Nick Ramsey, Jack Rogers, Robb Brumfield, Claire Thomas, Chloe St. Germain-Vermillion, Cham Mehaffey, Mary Mehaffey, Mitch Foret, Kathy Rhodes, William Matthews, Malise Prieto, Joan Garvey, Cathy DiSalvo
- 10. Light conditions (position of bird in relation to shade and to direction and amount of light): Full sun but photos appear dark because there is dense vegetation and trees shading the area.
- 11. Optical equipment (type, power, condition): Vortex Viper HD 10x42 Binoculars, Canon Rebel T6 with 150-600m Sigma telephoto lens.
- 12. Distance to bird(s): About 4 yds. when I first saw it.
- 13. Duration of observation: A solid 2 minutes at the feeder and hopping around the shrubs in my backyard on the first day that I saw it.
- 14. Habitat: Turk caps in a residential neighborhood.
- 15. Behavior of bird / circumstances of observation (flying, feeding, resting; include and stress habits used in identification; relate events surrounding observation): Vocalizing (chipping, rapid calling), feeding at my hummingbird feeder occasionally. Stay's in the same area all day. Responds to playback with "chetet" calls and staying perched in the area for a brief moment.

16. Description (include only what was actually seen, not what "should" have been seen; include if possible: total length/relative size compared to other familiar species, body bulk, shape, proportions, bill, eye, leg, and plumage characteristics. Stress features that separate it from similar species):

Long tail compared to other orioles. Big in comparison to other yellow birds. Curved bill.

- 17. Voice: chipping, chattering, "chetet" calls.
- 18. Similar species (include how they were eliminated by your observation): Orchard Oriole and Baltimore Oriole. They are the only other Orioles I've ever seen, so when I saw the Hooded Oriole I felt a bit suspicious. One because its shape was more slender looking, and because it's in its winter plumage/immature/female.
- 19. Photographs or tape recordings obtained? (by whom? attached?): Cham Mehaffey, Mary Mehaffey, Mitch Foret, Justin Bosler, William Bosler, Kathy Rhodes, Nick Ramsey, Jack Rogers, Claire Thomas, Chloe St. Germain-Vermillion, Robb Brumfield. Mitch Forets photos here: https://ebird.org/checklist/S162905372
- 20. Previous experience with this species: 0
- 21. Identification aids: (list books, illustrations, other birders, etc. used in identification):

Local birders, since I am from Connecticut and don't have any winter oriole experience.

- 22. Identification aids used when? At time of sighting, After observation
- 23. Are you positive of your identification? If not, explain: Yes
- 24. Date report completed: March 1st, 2024
- 25. May the LBRC have permission to display in whole or in part this report and accompanying photos on the LOS-LBRC website and LBRC Facebook page? Yes
- 26. If yes, may we include your name with the report? Yes









